

## Anti-STX17 Antibody Picoband® FITC Conjugated

Catalog Number: A06085-1-FITC

### About STX17

Enables SNAP receptor activity; SNARE binding activity; and protein phosphatase binding activity. Involved in several processes, including autophagosome membrane docking; endoplasmic reticulum to Golgi vesicle-mediated transport; and endoplasmic reticulum-Golgi intermediate compartment organization. Acts upstream of or within protein localization to phagophore assembly site. Located in several cellular components, including autophagosome membrane; endoplasmic reticulum-Golgi intermediate compartment; and mitochondria-associated endoplasmic reticulum membrane contact site. Part of SNARE complex.

### Overview

Product Name	Anti-STX17 Antibody Picoband® FITC Conjugated
Reactive Species	Human
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, IP, IF, IHC, ICC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na <sub>2</sub> HPO <sub>4</sub> , 0.02% NaN <sub>3</sub> .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P56962

### Technical Details

Immunogen	E.coli-derived human STX17 recombinant protein (Position: R12-H206).
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	FITC Excitation Wavelength: 495 nm Emission Wavelength: 525 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-STX17 Antibody - FITC

For Research Use Only. Not for use in diagnostic procedures.