

Anti-NAAA Antibody Picoband® Fluoro488 Conjugated

Catalog Number: A06195-1-Fluoro488

About NAAA

N-acyl ethanolamine-hydrolyzing acid amidase is an enzyme that in humans is encoded by the NAAA gene. Enables N-(long-chain-acyl)ethanolamine deacylase activity; N-acylsphingosine amidohydrolase activity; and fatty acid amide hydrolase activity. Involved in several processes, including N-acyl ethanolamine metabolic process; N-acyl phosphatidylethanolamine metabolic process; and sphingosine metabolic process. Located in lysosome. Is extrinsic component of membrane.

Overview

Product Name	Anti-NAAA Antibody Picoband® Fluoro488 Conjugated
Reactive Species	Human, Rat
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, Flow Cytometry, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% Na ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	Q02083

Technical Details

Immunogen	E.coli-derived human NAAA recombinant protein (Position: R35-L325).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	Fluoro488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-NAAA Antibody - Fluoro488

For Research Use Only. Not for use in diagnostic procedures.