

Anti-IL-18R Beta/IL-18RAP Antibody Picoband®

Catalog Number: A06457-2

About IL18RAP

Interleukin 18 receptor accessory protein, also known as IL18RAP and CDw218b (cluster of differentiation w218b), is a human gene. The protein encoded by this gene is an accessory subunit of the heterodimeric receptor for interleukin 18 (IL18), a proinflammatory cytokine involved in inducing cell-mediated immunity. This protein enhances the IL18-binding activity of the IL18 receptor and plays a role in signaling by IL18. Mutations in this gene are associated with Crohn's disease and inflammatory bowel disease, and susceptibility to celiac disease and leprosy. Alternatively spliced transcript variants of this gene have been described, but their full-length nature is not known.

Overview


Product Name	Anti-IL-18R Beta/IL-18RAP Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-IL-18R Beta/IL-18RAP Antibody Picoband® catalog # A06457-2. Tested in WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.005mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O95256

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human IL-18R Beta/IL-18RAP, which shares 84.4% amino acid (aa) sequence identity with both mouse and rat IL-18R Beta/IL-18RAP.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5ug/ml, Human

Anti-IL-18R Beta/IL-18RAP Antibody Picoband® (A06457-2) Images



Western blot analysis of IL-18R Beta/IL-18RAP using anti-IL-18R Beta/IL-18RAP antibody (A06457-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human HL-60 whole cell lysates, Lane 2: human A549 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-IL-18R Beta/IL-18RAP antigen affinity purified polyclonal antibody (Catalog # A06457-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for IL-18R Beta/IL-18RAP at approximately 72KD. The expected band size for IL-18R Beta/IL-18RAP is at 72KD.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-IL-18R Beta/IL-18RAP Antibody

For Research Use Only. Not for use in diagnostic procedures.