

Anti-KIR2.3 KCNJ4 Antibody

Catalog Number: A06605-1


Overview

Product Name	Anti-KIR2.3 KCNJ4 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-KIR2.3 KCNJ4 Antibody catalog # A06605-1. Tested in WB, ELISA, IHC applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, IHC, WB
Clonality	Polyclonal
Formulation	Liquid in PBS containing 50% glycerol, 0.5% stabilizing protein and 0.02% sodium azide. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P48050


Technical Details

Immunogen	The antiserum was produced against synthesized peptide derived from human KCNJ4. AA range:251-300
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	Immunogen affinity purified
Suggested Dilutions	WB 1:500-2000 IHC 1:50-300 ELISA 1:2000-20000


Anti-KIR2.3 KCNJ4 Antibody (A06605-1) Images



Western Blot analysis of 293 cells using KIR2.3 Polyclonal Antibody



Western blot analysis of KCNJ4 Antibody. The lane on the right is blocked with the KCNJ4 peptide.



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200 (4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-KIR2.3 KCNJ4 Antibody

For Research Use Only. Not for use in diagnostic procedures.