BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

# Anti-Nexilin NEXN Antibody

**Catalog Number:** A06648

## Submit a product review to Biocompare.com







Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

## Overview


| | |
|---|---|
| Product Name | Anti-Nexilin NEXN Antibody |
| Reactive Species | Human, Mouse, Rat |
| Description | Boster Bio Anti-Nexilin NEXN Antibody catalog # A06648. Tested in WB,ICC/IF applications. This antibody reacts with Human,Mouse,Rat. |
| Application | IF, ICC, WB |
| Clonality | Polyclonal |
| Formulation | Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2 |
| Storage Instructions | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles. |
| Host | Rabbit |
| Uniprot ID | Q0ZGT2 |

## Technical Details


| | |
|---|---|
| Immunogen | Recombinant fusion protein of human NEXN(NP_653174.3). |
| Cross Reactivity | No cross reactivity with other proteins. |
| Isotype | IgG |
| Form | Liquid |
| Concentration | 1 mg/ml |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE). |
| Suggested Dilutions | WB: 1:500-1:2000<br>ICC/IF: 1:50-1:200 |

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

# Anti-Nexilin NEXN Antibody (A06648) Images




Western blot analysis of extracts of various cell lines, using NEXN antibody at 1:3000 dilution.Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.Lysates/proteins: 25ug per lane.Blocking buffer: 3% nonfat dry milk in TBST.Detection: ECL Basic Kit .Exposure time: 90s.



Immunofluorescence analysis of U-2 OS cells using NEXN antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

Anti-Nexilin NEXN Antibody

For Research Use Only. Not for use in diagnostic procedures.