

## Anti-TTL Antibody Picoband® APC Conjugated

Catalog Number: A06710-APC

### About TTL

TTL is a cytosolic enzyme involved in the posttranslational modification of alpha-tubulin (see MIM 602529). Alpha-tubulin within assembled microtubules is detyrosinated over time at the C terminus. After microtubule disassembly, TTL restores the tyrosine residues and consequently participates in a cycle of tubulin detyrosination and tyrosination (Erck et al., 2003 [PubMed 14571137]).

### Overview

Product Name	Anti-TTL Antibody Picoband® APC Conjugated
Reactive Species	Mouse, Rat
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na <sub>2</sub> HPO <sub>4</sub> , 0.02% NaN <sub>3</sub> .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	Q8NG68

### Technical Details

Immunogen	E.coli-derived human TTL recombinant protein (Position: N104-L377). Human TTL shares 95.3% and 95.6% amino acid (aa) sequence identity with mouse and rat TTL, respectively.
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	APC Excitation Wavelength: 633-647 nm Emission Wavelength: 660 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-TTL Antibody - APC

For Research Use Only. Not for use in diagnostic procedures.