

Anti-Beta Sarcoglycan/SGCB Antibody Picoband®

Catalog Number: A07064-2

About SGCB

This gene encodes a member of the sarcoglycan family. Sarcoglycans are transmembrane components in the dystrophin-glycoprotein complex which help stabilize the muscle fiber membranes and link the muscle cytoskeleton to the extracellular matrix. Mutations in this gene have been associated with limb-girdle muscular dystrophy.


Overview

Product Name	Anti-Beta Sarcoglycan/SGCB Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Beta Sarcoglycan/SGCB Antibody Picoband® catalog # A07064-2. Tested in WB, ELISA applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q16585

Technical Details

Immunogen	E.coli-derived human Beta Sarcoglycan/SGCB recombinant protein (Position: E101-H318). Human Beta Sarcoglycan/SGCB shares 95.4% amino acid (aa) sequence identity with mouse Beta Sarcoglycan/SGCB.
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Human, Mouse, Rat ELISA, 0.1-0.5 ug/ml

Anti-Beta Sarcoglycan/SGCB Antibody Picoband® (A07064-2) Images



Western blot analysis of Beta Sarcoglycan/SGCB using anti-Beta Sarcoglycan/SGCB antibody (A07064-2). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human SIHA whole cell lysates, Lane 2: human HT1080 whole cell lysates, Lane 3: rat heart tissue lysates, Lane 4: mouse heart tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Beta Sarcoglycan/SGCB antigen affinity purified polyclonal antibody (A07064-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for Beta Sarcoglycan/SGCB at approximately 45 kDa. The expected band size for Beta Sarcoglycan/SGCB is at 35 kDa.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Beta Sarcoglycan/SGCB Antibody

For Research Use Only. Not for use in diagnostic procedures.