

Anti-UBAP2L Antibody Picoband®

Catalog Number: A07183-3

About UBAP2L

Ubiquitin-associated protein 2-like is a protein that in humans is encoded by the UBAP2L gene. Ubiquitin-associated protein 2-like (UBAP2L), which contains a ubiquitin-associated (UBA) domain near its N-terminus, has been indicated in the pathogenesis of several human cancers, including multiple myeloma, hepatocellular carcinoma, and malignant ovarian tumors. UBAP2L is closely associated with tumor growth and metastasis. UBAP2L has a key role in glioma cell growth and may act as an oncogene to promote malignant glioma development. UBAP2L also plays an important role in breast cancer cell proliferation and might serve as a potential target for breast cancer treatment. UBAP2L plays a critical role in the maintenance of the metastatic ability of hepatocellular carcinoma (HCC) cells via SNAIL1 Regulation and is predictive of a poor clinical outcome.

Overview


Product Name	Anti-UBAP2L Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-UBAP2L Antibody Picoband® catalog # A07183-3. Tested in ELISA, Flow Cytometry, IF, ICC, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, IF, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q14157

Technical Details


Immunogen	E.coli-derived human UBAP2L recombinant protein (Position: A271-S439).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for ICC.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.25 ug/ml, Human Immunocytochemistry/Immunofluorescence, 5 ug/ml, Human Flow Cytometry (Fixed), 1-3 ug/1x10 ⁶ cells, Human ELISA, 0.1-0.5 ug/ml, -


Anti-UBAP2L Antibody Picoband® (A07183-3) Images



Western blot analysis of UBAP2L using anti-UBAP2L antibody (A07183-3). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Jurkat whole cell lysates, Lane 2: human K562 whole cell lysates, Lane 3: human PC-3 whole cell lysates, Lane 4: human Hela whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-UBAP2L antigen affinity purified polyclonal antibody (Catalog # A07183-3) at 0.25 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for UBAP2L at approximately 150 kDa. The expected band size for UBAP2L is at 150 kDa.



IF analysis of UBAP2L using anti-UBAP2L antibody (A07183-3). UBAP2L was detected in an immunocytochemical section of T-47D cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-UBAP2L Antibody (A07183-3) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



Flow Cytometry analysis of THP-1 cells using anti-UBAP2L antibody (A07183-3). Overlay histogram showing THP-1 cells stained with A07183-3 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-UBAP2L Antibody (A07183-3, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-UBAP2L Antibody

For Research Use Only. Not for use in diagnostic procedures.