

Anti-CIR Antibody (N-term)

Catalog Number: A07268

About CIR1

CIR may modulate splice site selection during alternative splicing of pre-mRNAs. It regulates transcription and acts as corepressor for RBPSUH. It recruits RBPSUH to the Sin3-histone deacetylase complex (HDAC) and is required for RBPSUH-mediated repression of transcription.


Overview

Product Name	Anti-CIR Antibody (N-term)
Reactive Species	Human
Description	Boster Bio Anti-CIR Antibody (N-term) (Catalog # A07268). Tested in WB application(s). This antibody reacts with Human.
Application	WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q86X95

Technical Details

Immunogen	This CIR antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 60-88 amino acids from the N-terminal region of human CIR.
Predicted Reactive Species	Bovine, Mouse, Pig
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	WB: 1:1000

Anti-CIR Antibody (N-term) (A07268) Images



CIR Antibody (N-term) western blot analysis in THP-1 cell line, human placenta, rat heart and skeletal muscle lysates (35ug/lane). This demonstrates the CIR antibody detected the CIR protein (arrow).

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CIR Antibody (N-term)

For Research Use Only. Not for use in diagnostic procedures.