

Anti-ICE2 Antibody Picoband®

Catalog Number: A07414-1

About ICE2

This gene encodes a protein component of the little elongation complex (LEC), which plays a role in small nuclear RNA (snRNA) transcription. The LEC regulates snRNA transcription by enhancing both RNA Polymerase II occupancy and transcriptional elongation. The encoded protein and other LEC components have been shown to localize to Cajal bodies, which are sites of ribonucleoprotein (RNP) complex assembly. Pseudogenes of this gene have been identified on chromosomes 3 and 4.

Submit a product review to Biocompare.com

[Submit a review of this product](#) to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Overview


Product Name	Anti-ICE2 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-ICE2 Antibody Picoband® catalog # A07414-1. Tested in WB, ELISA applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q659A1

Technical Details

Immunogen	E.coli-derived human ICE2 recombinant protein (Position: I522-D533).
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Human, Mouse, Rat ELISA, 0.1-0.5 ug/ml

Anti-ICE2 Antibody Picoband® (A07414-1) Images



Western blot analysis of ICE2 using anti-ICE2 antibody (A07414-1). Electrophoresis was performed on a 8% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human A549 whole cell lysates, Lane 2: human U251 whole cell lysates, Lane 3: human K562 whole cell lysates, Lane 4: human SH-SY5Y whole cell lysates, Lane 5: rat RH35 whole cell lysates, Lane 6: rat C6 whole cell lysates, Lane 7: mouse HEPA1-6 whole cell lysates, Lane 8: mouse Neuro-2a whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ICE2 antigen affinity purified polyclonal antibody (A07414-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for ICE2 at approximately 110 kDa. The expected band size for ICE2 is at 110 kDa.

Anti-ICE2 Antibody

For Research Use Only. Not for use in diagnostic procedures.