

Anti-SLC26A9 Antibody Picoband®

Catalog Number: A07581-1

About SLC26A9

The members of the SLC26 family of transporters, such as SLC26A9, are well conserved across species and mediate the electroneutral exchange of chloride for bicarbonate or sulfate across the plasma membrane.

Overview

Product Name	Anti-SLC26A9 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-SLC26A9 Antibody Picoband® catalog # A07581-1. Tested in ELISA, WB, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q7LBE3


Technical Details

Immunogen	E.coli-derived human SLC26A9 recombinant protein (Position: K27-D763). Human SLC26A9 shares 89.7% amino acid (aa) sequence identity with mouse SLC26A9.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.
Purification	Immunogen affinity purified.


Suggested Dilutions

Western blot, 0.25-0.5 ug/ml, Human, Mouse, Rat
Flow Cytometry (Fixed), 1-3 ug/ 1×10^6 cells, Human
ELISA, 0.1-0.5 ug/ml, -

Anti-SLC26A9 Antibody Picoband® (A07581-1) Images



Western blot analysis of SLC26A9 using anti-SLC26A9 antibody (A07581-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates, Lane 2: rat stomach tissue lysates, Lane 3: rat lung tissue lysates, Lane 4: rat C6 whole cell lysates, Lane 5: mouse stomach tissue lysates, Lane 6: mouse lung tissue lysates, Lane 7: mouse MFC whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SLC26A9 antigen affinity purified polyclonal antibody (Catalog # A07581-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for SLC26A9 at approximately 90 kDa. The expected band size for SLC26A9 is at 87 kDa.



Flow Cytometry analysis of HepG2 cells using anti-SLC26A9 antibody (A07581-1). Overlay histogram showing HepG2 cells stained with A07581-1 (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-SLC26A9 Antibody (A07581-1, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (A07581-1, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-SLC26A9 Antibody

For Research Use Only. Not for use in diagnostic procedures.