

## Anti-GMIP Antibody

Catalog Number: A07655-1

### About GMIP

GMIP (Gem-interacting protein) is a 970 amino acid protein that stimulates the GTPase activity of RhoA in vitro and in vivo. GMIP interacts with Gem through its N-terminus and has a Rho GTPase-activating protein domain at its C-terminus. The Rho family of GTP-binding proteins plays a role in the development of neuronal structure. GMIP is able to inhibit RhoA function, leading to Actin cytoskeletal reorganization in vivo.

### Overview


Product Name	Anti-GMIP Antibody
Reactive Species	Human
Description	Boster Bio Anti-GMIP Antibody catalog # A07655-1. Tested in WB, IHC, ICC, IF, IP, ELISA applications. This antibody reacts with Human.
Application	ELISA, IP, IF, IHC, ICC, WB
Clonality	Polyclonal
Formulation	500 ug/ml antibody with PBS, 0.02% NaN <sub>3</sub> , 1 mg stabilizing protein and 50% glycerol This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	12 months from date of receipt $-20^{\circ}\text{C}$ as supplied. 6 months $2$ to $8^{\circ}\text{C}$ after reconstitution. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q9P107

### Technical Details

Immunogen	E.coli-derived human GMIP recombinant protein (Position: A3-R89).
Form	Liquid
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 1:500-2000 Immunohistochemistry, 1:50-400 Immunocytochemistry/Immunofluorescence, 1:50-400 ImmunoPrecipitation, 1:250-300 ELISA, 1:100-1000



## Anti-GMIP Antibody (A07655-1) Images



Western blot analysis of GMIP using anti-GMIP antibody (A07655-1). Electrophoresis was performed on a 8% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human THP-1 whole cell lysates, Lane 2: human RT4 whole cell lysates, Lane 3: human Jurkat whole cell lysates, Lane 4: human HEL whole cell lysates, Lane 5: mouse thymus tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-GMIP antigen affinity purified polyclonal antibody (A07655-1) at 1:1000 overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for GMIP at approximately 130 kDa. The expected band size for GMIP is at 107 kDa.

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-GMIP Antibody

For Research Use Only. Not for use in diagnostic procedures.