

Anti-TAS1R1 Antibody Picoband®

Catalog Number: A07896-1

About TAS1R1

Predicted to enable G protein-coupled receptor activity and taste receptor activity. Predicted to be involved in sensory perception of umami taste. Predicted to be active in plasma membrane. Is expressed in tongue anterior part. Orthologous to human TAS1R1 (taste 1 receptor member 1).


Overview

Product Name	Anti-TAS1R1 Antibody Picoband®
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-TAS1R1 Antibody Picoband® catalog # A07896-1. Tested in WB, ELISA applications. This antibody reacts with Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q99PG6

Technical Details

Immunogen	E.coli-derived mouse TAS1R1 recombinant protein (Position: E102-T842). Mouse TAS1R1 shares 72.9% and 88.8% amino acid (aa) sequence identity with human and rat TAS1R1, respectively.
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Mouse, Rat ELISA, 0.1-0.5 ug/ml

Anti-TAS1R1 Antibody Picoband® (A07896-1) Images



Western blot analysis of TAS1R1 using anti-TAS1R1 antibody (A07896-1). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat testis tissue lysates, Lane 2: rat C6 whole cell lysates, Lane 3: mouse testis tissue lysates, Lane 4: mouse brain tissue lysates, Lane 5: mouse Neuro-2a whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TAS1R1 antigen affinity purified polyclonal antibody (A07896-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for TAS1R1 at approximately 140 kDa. The expected band size for TAS1R1 is at 93 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TAS1R1 Antibody

For Research Use Only. Not for use in diagnostic procedures.