

Anti-DPH1 Antibody Picoband®

Catalog Number: A08287-1

About DPH1

The protein encoded by this gene is an enzyme involved in the biosynthesis of diphthamide, a modified histidine found only in elongation factor-2 (EEF2). Diphthamide residues in EEF2 are targeted for ADP-ribosylation by diphtheria toxin and Pseudomonas exotoxin A. Defects in this gene have been associated with both ovarian cancer and autosomal recessive intellectual disability with short stature, craniofacial, and ectodermal anomalies.


Overview

Product Name	Anti-DPH1 Antibody Picoband®
Reactive Species	Human, Rat
Description	Boster Bio Anti-DPH1 Antibody Picoband® catalog # A08287-1. Tested in WB, ELISA applications. This antibody reacts with Human, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q9BZG8

Technical Details

Immunogen	E.coli-derived human DPH1 recombinant protein (Position: P50-P438).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Human, Rat ELISA, 0.1-0.5 ug/ml

Anti-DPH1 Antibody Picoband® (A08287-1) Images



Western blot analysis of DPH1 using anti-DPH1 antibody (A08287-1). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Hela whole cell lysates, Lane 2: human 293T whole cell lysates, Lane 3: rat kidney tissue lysates, Lane 4: rat liver tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-DPH1 antigen affinity purified polyclonal antibody (A08287-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for DPH1 at approximately 60 kDa. The expected band size for DPH1 is at 48 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-DPH1 Antibody

For Research Use Only. Not for use in diagnostic procedures.