

## Anti-UFL1 Antibody

Catalog Number: A08383-1

### About UFL1

Enables UFM1 ligase activity and protein kinase binding activity. Involved in several processes, including positive regulation of reticulophagy; regulation of intracellular signal transduction; and regulation of primary metabolic process. Acts upstream of or within several processes, including positive regulation of cell population proliferation; regulation of proteasomal ubiquitin-dependent protein catabolic process; and response to endoplasmic reticulum stress. Located in cytoplasm; nucleus; and site of double-strand break. Part of protein-containing complex. Is active in endoplasmic reticulum membrane and mitochondrial outer membrane.

### Overview


Product Name	Anti-UFL1 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-UFL1 Antibody catalog # A08383-1. Tested in WB, IHC, ELISA applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, IHC, WB
Clonality	Polyclonal
Formulation	500 ug/ml antibody with PBS, 0.02% NaN <sub>3</sub> , 1 mg stabilizing protein and 50% glycerol This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	12 months from date of receipt at -20°C as supplied. 6 months at 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	O94874

### Technical Details


Immunogen	E.coli-derived human UFL1 recombinant protein (Position: 3-347).
Form	Liquid
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 1:500-2000 Immunohistochemistry, 1:50-400 ELISA, 1:100-1000



## Anti-UFL1 Antibody (A08383-1) Images



Western blot analysis of UFL1 using anti-UFL1 antibody (A08383-1). Electrophoresis was performed on a 8% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates, Lane 2: human K562 whole cell lysates, Lane 4: rat PC-12 whole cell lysates, Lane 5: mouse Neuro-2a whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-UFL1 antigen affinity purified polyclonal antibody (A08383-1) at 1:1000 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for UFL1 at approximately 100 kDa. The expected band size for UFL1 is at 90 kDa.



IHC analysis of UFL1 using anti-UFL1 antibody (A08383-1). UFL1 was detected in a paraffin-embedded section of human thyroid cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:100 rabbit anti-UFL1 Antibody (A08383-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

### Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-UFL1 Antibody

For Research Use Only. Not for use in diagnostic procedures.