

Anti-LAPTM4B antibody

Catalog Number: A08504


Overview

| | |
|----------------------|--|
| Product Name | Anti-LAPTM4B antibody |
| Reactive Species | Human, Mouse, Rat |
| Description | Boster Bio Anti-LAPTM4B antibody catalog # A08504. Tested in WB, IHC, ELISA applications. This antibody reacts with Human, Mouse, Rat. |
| Application | ELISA, IHC, WB |
| Clonality | Polyclonal |
| Formulation | 500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg stabilizing protein and 50% glycerol This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required. |
| Storage Instructions | 12 months from date of receipt at -20°C as supplied. 6 months at 2 to 8°C after reconstitution. Avoid repeated freezing and thawing. |
| Host | Rabbit |
| Uniprot ID | Q86VI4 |


Technical Details

| | |
|---------------------|---|
| Immunogen | E.coli-derived human LAPTM4B recombinant protein (Position:) |
| Form | Liquid |
| Concentration | 500 ug/ml |
| Purification | Immunogen affinity purified. |
| Suggested Dilutions | Western blot, 1:500-2000 Immunohistochemistry, 1:50-400 ELISA, 1:100-1000 |


Anti-LAPTM4B antibody (A08504) Images



IHC analysis of LAPTM4B using anti-LAPTM4B antibody (A08504). LAPTM4B was detected in a paraffin-embedded section of human thyroid cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1: 200 rabbit anti-LAPTM4B Antibody (A08504) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of LAPTM4B using anti-LAPTM4B antibody (A08504). LAPTM4B was detected in a paraffin-embedded section of human thyroid cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1: 200 rabbit anti-LAPTM4B Antibody (A08504) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of LAPTM4B using anti-LAPTM4B antibody (A08504). LAPTM4B was detected in a paraffin-embedded section of rat ovary tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1: 200 rabbit anti-LAPTM4B Antibody (A08504) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-LAPTM4B antibody

For Research Use Only. Not for use in diagnostic procedures.