

Anti-PSMD11 (R310) Antibody

Catalog Number: A08940

About PSMD11

Has RNA-dependent ATPase activity. Represses the transcriptional activity of nuclear receptors.

Scherer S.E., Nature 440:346-351(2006) [PubMed: 16541075].

Rajendran R.R., J. Biol. Chem. 278:4628-4638(2003) [PubMed: 12466272].

Scherl A., Mol. Biol. Cell 13:4100-4109(2002) [PubMed: 12429849].

Overview

| Product Name | Anti-PSMD11 (R310) Antibody |
|----------------------|--------------------------------------------------------------------------------------------------------------------------------------------|
| Reactive Species | Human, Mouse, Rat |
| Description | Boster Bio Anti-PSMD11 (R310) Antibody catalog # A08940. Tested in WB,IHC applications. This antibody reacts with Human,Mouse,Rat. |
| Application | IHC, WB |
| Clonality | Polyclonal |
| Formulation | Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2 |
| Storage Instructions | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles. |
| Host | Rabbit |
| Uniprot ID | O00231 |

Technical Details

| Immunogen | Synthesized peptide derived from human NU205 protein. |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Predicted Reactive Species | Chimpanzee, Drosophila, Macaque |
| Isotype | IgG |
| Form | Liquid |
| Concentration | 1 mg/ml |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE). |
| Suggested Dilutions | Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. |



BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

| optimal Some P Boster WB: 1:5 | pected range of concentration is unknown, a pilot test should be conducted to decide the dilution ratio for your samples. ubMed article(s) citing the expression level of this target are as follows: 3io's internal QC testing used: 00-1:1000 50-1:200 |
|----------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|----------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-PSMD11 (R310) Antibody