

Anti-Carbonic Anhydrase 13/CA13 Antibody Picoband®

Catalog Number: A09186-3

About CA13

Carbonic anhydrase 13 is a protein that in humans is encoded by the CA13 gene. Carbonic anhydrases (CAs) are a family of zinc metalloenzymes that catalyze the interconversion between carbon dioxide and water and the dissociated ions of carbonic acid (i.e. bicarbonate and hydrogen ions).

Overview


Product Name	Anti-Carbonic Anhydrase 13/CA13 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Carbonic Anhydrase 13/CA13 Antibody Picoband® catalog # A09186-3. Tested in Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	Flow Cytometry, IF, IHC, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.01mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q8N1Q1

Technical Details


Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human Carbonic Anhydrase 13/CA13, which shares 85% and 55% amino acid (aa) sequence identity with mouse and rat Carbonic Anhydrase 13/CA13, respectively.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P) and ICC.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5ug/ml, Human, Mouse, Rat Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Human Immunocytochemistry/Immunofluorescence, 5ug/ml, Human Flow Cytometry (Fixed), 1-3ug/1x10 ⁶ cells, Human


Anti-Carbonic Anhydrase 13/CA13 Antibody Picoband® (A09186-3) Images




Western blot analysis of Carbonic Anhydrase 13/CA13 using anti-Carbonic Anhydrase 13/CA13 antibody (A09186-3). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates, Lane 2: human CACO-2 whole cell lysates, Lane 3: human U87 whole cell lysates, Lane 4: human K562 whole cell lysates, Lane 5: human Hela whole cell lysates, Lane 6: rat intestines tissue lysates, Lane 7: mouse NTH/3T3 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Carbonic Anhydrase 13/CA13 antigen affinity purified polyclonal antibody (Catalog # A09186-3) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Carbonic Anhydrase 13/CA13 at approximately 29KD. The expected band size for Carbonic Anhydrase 13/CA13 is at 29KD.




IHC analysis of Carbonic Anhydrase 13/CA13 using anti-Carbonic Anhydrase 13/CA13 antibody (A09186-3). Carbonic Anhydrase 13/CA13 was detected in paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-Carbonic Anhydrase 13/CA13 Antibody (A09186-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.




IHC analysis of Carbonic Anhydrase 13/CA13 using anti-Carbonic Anhydrase 13/CA13 antibody (A09186-3). Carbonic Anhydrase 13/CA13 was detected in paraffin-embedded section of human melanoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-Carbonic Anhydrase 13/CA13 Antibody (A09186-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.




IHC analysis of Carbonic Anhydrase 13/CA13 using anti-Carbonic Anhydrase 13/CA13 antibody (A09186-3). Carbonic Anhydrase 13/CA13 was detected in paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-Carbonic Anhydrase 13/CA13 Antibody (A09186-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.



IHC analysis of Carbonic Anhydrase 13/CA13 using anti-Carbonic Anhydrase 13/CA13 antibody (A09186-3). Carbonic Anhydrase 13/CA13 was detected in paraffin-embedded section of human ovarian cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-Carbonic Anhydrase 13/CA13 Antibody (A09186-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.



IF analysis of Carbonic Anhydrase 13/CA13 using anti-Carbonic Anhydrase 13/CA13 antibody (A09186-3). Carbonic Anhydrase 13/CA13 was detected in immunocytochemical section of Hep cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5ug/mL rabbit anti-Carbonic Anhydrase 13/CA13 Antibody (A09186-3) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



Flow Cytometry analysis of CACO-2 cells using anti-Carbonic Anhydrase 13/CA13 antibody (A09186-3). Overlay histogram showing CACO-2 cells stained with A09186-3 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Carbonic Anhydrase 13/CA13 Antibody (A09186-3, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with

primary antibody and secondary antibody (Red line) was used as a blank control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Carbonic Anhydrase 13/CA13 Antibody

For Research Use Only. Not for use in diagnostic procedures.