

Anti-CD72 Antibody Picoband®

Catalog Number: A09292-1

About CD72

CD72 (Cluster of Differentiation 72), also known in murine biology as Lyb-2, is a protein active in the immune system of animals. This gene is mapped to 9p13.3. It consists of two identical halves, each of about 39-43 kD, and is a C-type lectin. Its primary locus of expression is B-cells, where it appears to mediate aspects of B-cell - T-cell interaction. It is a ligand for CD5.

Overview


Product Name	Anti-CD72 Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-CD72 Antibody Picoband® catalog # A09292-1. Tested in ELISA, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Na ₃ N.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P21854

Technical Details

Immunogen	E. coli-derived human CD72 recombinant protein (Position: R117-D359).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml

	ELISA, 0.1-0.5ug/ml
--	---------------------

Anti-CD72 Antibody Picoband® (A09292-1) Images



Western blot analysis of CD72 using anti-CD72 antibody (A09292-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human A375 whole cell lysate. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CD72 antigen affinity purified polyclonal antibody (Catalog # A09292-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CD72 at approximately 45KD. The expected band size for CD72 is at 40KD.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CD72 Antibody

For Research Use Only. Not for use in diagnostic procedures.