

## Anti-LAPTM5 Antibody (N-term)

Catalog Number: A09673

### About LAPTM5

RNA-binding protein that plays a role in the regulation of alternative splicing and influences mRNA splice site selection and exon inclusion. Its phosphorylation by FYN inhibits its ability to regulate splice site selection. Induces an increased concentration-dependent incorporation of exon in CD44 pre-mRNA by direct binding to purine-rich exonic enhancer. May function as an adapter protein for Src kinases during mitosis. Binds both poly(A) and poly(U) homopolymers. Phosphorylation by PTK6 inhibits its RNA-binding ability (By similarity).


### Overview

Product Name	Anti-LAPTM5 Antibody (N-term)
Reactive Species	Human
Description	Boster Bio Anti-LAPTM5 Antibody (N-term) (Catalog # A09673). Tested in WB, Flow Cytometry, IHC-P application(s). This antibody reacts with Human.
Application	Flow Cytometry, IHC-P, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q13571


### Technical Details

Immunogen	This LAPTM5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 23-49 amino acids from the N-terminal region of human LAPTM5.
Predicted Reactive Species	Bovine
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	WB: 1:1000 IHC-P: 1:50-1:100 FC: 1:10-1:50


## Anti-LAPTM5 Antibody (N-term) (A09673) Images



All lanes : Anti-LAPTM5 Antibody (N-term) at 1:1000 dilution  
Lane 1: U266B1 whole cell lysate  
Lane 2: Jurkat whole cell lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 30 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.



LAPTM5 antibody (N-term) immunohistochemistry analysis in formalin fixed and paraffin embedded human lymph node followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the LAPTM5 antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



LAPTM5 Antibody (N-term) flow cytometric analysis of NCI-H460 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

**Submit a product review to Biocompare.com**

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-LAPTM5 Antibody (N-term)

For Research Use Only. Not for use in diagnostic procedures.