

Anti-BEX3 Antibody Picoband®

Catalog Number: A09675-2

About BEX3

Protein BEX3 is a protein that in humans is encoded by the NGFRAP1 gene. NGFRAP1 (nerve growth factor receptor associated protein 1), also called BEX3 (brain-expressed X-linked protein 3), NADE (p75NTR-associated death executor) or FAP-1 (Fas-associated phosphatase-1), is a 24 kDa member of the BEX protein family. A nuclear export signal (aa 77-87) allows NGFRAP1 to export bound p75NTR. It is widely expressed, but highest in ovarian granulosa cells, testis, prostate, seminal vesicle and liver. Except for a 13-19 amino acid (aa) rodent Arg/His region at aa 35 that is not found in humans, human NGFRAP1 shares 94% aa identity with mouse and rat NGFRAP1. In humans, a 101 aa isoform with an alternative start site at aa 11 has been described.

Overview


Product Name	Anti-BEX3 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-BEX3 Antibody Picoband® catalog # A09675-2. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q00994

Technical Details


Immunogen	E.coli-derived human BEX3 recombinant protein (Position: Q39-H103).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Human, Mouse, Rat Flow Cytometry (Fixed), 1-3 ug/1x10 ⁶ cells, Human ELISA, 0.1-0.5 ug/ml, -

Anti-BEX3 Antibody Picoband® (A09675-2) Images



Western blot analysis of BEX3 using anti-BEX3 antibody (A09675-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HCCT tissue lysates, Lane 2: human HCCP tissue lysates, Lane 3: human MCF-7 whole cell lysates, Lane 4: human Hela whole cell lysates, Lane 5: rat liver tissue lysates, Lane 6: rat L6 whole cell lysates, Lane 7: mouse liver tissue lysates, Lane 8: mouse C2C12 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-BEX3 antigen affinity purified polyclonal antibody (Catalog # A09675-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for BEX3 at approximately 28 kDa. The expected band size for BEX3 is at 28 kDa.



Flow Cytometry analysis of MCF-7 cells using anti-BEX3 antibody (A09675-2). Overlay histogram showing MCF-7 cells stained with A09675-2 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-BEX3 Antibody (A09675-2, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-BEX3 Antibody

For Research Use Only. Not for use in diagnostic procedures.