

Anti-DNAJA4 Antibody Picoband®

Catalog Number: A10527-1

About DNAJA4

Enables protein-folding chaperone binding activity and unfolded protein binding activity. Involved in several processes, including negative regulation of endothelial cell migration; negative regulation of inclusion body assembly; and protein refolding. Located in cytosol and membrane.

Submit a product review to Biocompare.com

[Submit a review of this product](#) to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Overview

Product Name	Anti-DNAJA4 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-DNAJA4 Antibody Picoband® catalog # A10527-1. Tested in WB, ELISA applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q8WW22


Technical Details

Immunogen	E.coli-derived human DNAJA4 recombinant protein (Position: K275-A390).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.

Suggested Dilutions

Western blot, 0.25-0.5 ug/ml, Human, Mouse, Rat
ELISA, 0.1-0.5 ug/ml

Anti-DNAJA4 Antibody Picoband® (A10527-1) Images



Western blot analysis of DNAJA4 using anti-DNAJA4 antibody (A10527-1). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human SiHa whole cell lysates, Lane 2: human THP-1 whole cell lysates, Lane 3: human A549 whole cell lysates, Lane 4: rat testis tissue lysates, Lane 5: rat heart tissue lysates, Lane 6: mouse testis tissue lysates, Lane 7: mouse heart tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-DNAJA4 antigen affinity purified polyclonal antibody (A10527-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for DNAJA4 at approximately 45 kDa. The expected band size for DNAJA4 is at 45 kDa.

Anti-DNAJA4 Antibody

For Research Use Only. Not for use in diagnostic procedures.