BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-PI 3-Kinase C2 gamma PIK3C2G Antibody

Catalog Number: A10773

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview


Product Name	Anti-PI 3-Kinase C2 gamma PIK3C2G Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-PI 3-Kinase C2 gamma PIK3C2G Antibody catalog # A10773. Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Liquid in PBS containing 50% glycerol, 0.5% stabilizing protein and 0.02% sodium azide. This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O75747

Technical Details


Immunogen	Synthesized peptide derived from the N-terminal region of human PI 3-Kinase C2gamma.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	Immunogen affinity purified
Suggested Dilutions	WB 1:500-2000 IHC 1:50-300

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-PI 3-Kinase C2 gamma PIK3C2G Antibody (A10773) Images



Western blot analysis of A549 using PI 3-Kinase C2gamma antibody. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200 (4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).

1 Publications Citing This Product

1. PubMed ID: 10.1016/j.envpol.2021.116556, Mechanisms underlying reproductive toxicity induced by nickel nanoparticles identified by comprehensive gene expression analysis in GC-1 spg cells

Visit bosterbio.com/anti-pi-3-kinase-c2-gamma-antibody-a10773-boster.html to see all 1 publications.

Anti-PI 3-Kinase C2 gamma PIK3C2G Antibody

For Research Use Only. Not for use in diagnostic procedures.