

Anti-MTCL1 Antibody Picoband® FITC Conjugated

Catalog Number: A10954-1-FITC

About MTCL1

MTCL1 (microtubule cross-linking factor 1), also known as CCDC165 (Coiledcoil domain-containing protein 165), SOGA2 or MTCL1, is a 1,905 amino acid protein that localizes to the cell membrane, cytoplasm and cytoskeleton. MTCL1 is a microtubule-associated factor that plays a role in regulating polarization and microtubule dynamics as well as the development and maintenance of non-centrosomal microtubule bundles. MTCL1 is encoded by a gene that maps to chromosome 18 and is expressed as four isoforms due to alternative splicing events. Chromosome 18 encodes over 300 genes and contains about 76 million bases. Trisomy 18, or Edwards syndrome, is the second most common trisomy after Downs syndrome. Symptoms of Edwards syndrome include low birth weight, a variety of physical development defects, heart deformations and breathing difficulty. Translocation between chromosome 18 and 14 is the most common translocation in cancers, and occurs in follicular lymphomas. Niemann-Pick disease, hereditary hemorrhagic telangiectasia and erythropoietic protoporphyria are associated with chromosome 18. The TGFbeta modulators, Smad2, Smad4 and Smad7 are encoded by chromosome 18.

Overview

Product Name	Anti-MTCL1 Antibody Picoband® FITC Conjugated
Reactive Species	Human, Mouse, Rat
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, Flow Cytometry, IHC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% Na ₃ N.
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	Q9Y4B5

Technical Details

Immunogen	E.coli-derived human MTCL1 recombinant protein (Position: K884-D1522).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.

Conjugate	FITC Excitation Wavelength: 495 nm Emission Wavelength: 525 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-MTCL1 Antibody - FITC

For Research Use Only. Not for use in diagnostic procedures.