

Anti-NPAP1 Antibody Picoband®

Catalog Number: A11015

About NPAP1

This intronless retrogene is located in the Prader-Willi syndrome region on chromosome 15. This gene exhibits tissue-specific imprinting. Expression in adult testis and brain is biallelic, while expression in fetal brain is monoallelic and only from the paternal chromosome. The encoded protein is associated with the nuclear pore complex.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Overview


Product Name	Anti-NPAP1 Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-NPAP1 Antibody Picoband® catalog # A11015. Tested in WB, ELISA applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q9NZP6

Technical Details

Immunogen	E.coli-derived human NPAP1 recombinant protein (Position: A153-P1156).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Human ELISA, 0.1-0.5 ug/ml

Anti-NPAP1 Antibody Picoband® (A11015) Images



Western blot analysis of NPAP1 using anti-NPAP1 antibody (A11015). Electrophoresis was performed on a 8% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human U2OS whole cell lysates, Lane 2: human RT4 whole cell lysates, Lane 3: human K562 whole cell lysates, Lane 4: human Caco-2 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-NPAP1 antigen affinity purified polyclonal antibody (A11015) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for NPAP1 at approximately 121 kDa. The expected band size for NPAP1 is at 121 kDa.

Anti-NPAP1 Antibody

For Research Use Only. Not for use in diagnostic procedures.