

Anti-DCHS2 Antibody Picoband®

Catalog Number: A11158

About DCHS2

This gene encodes a large protein that contains many cadherin domains and likely functions in cell adhesion. Genome-wide association studies suggest that this gene may be important in Alzheimer's disease, compressive strength index, and appendicular lean mass.

Submit a product review to Biocompare.com

[Submit a review of this product](#) to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Overview

Product Name	Anti-DCHS2 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-DCHS2 Antibody Picoband® catalog # A11158. Tested in WB, ELISA applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q6V1P9


Technical Details

Immunogen	E.coli-derived human DCHS2 recombinant protein (Position: I858-E2983).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.

Suggested Dilutions

Western blot, 0.25-0.5 ug/ml, Human, Mouse, Rat
ELISA, 0.1-0.5 ug/ml

Anti-DCHS2 Antibody Picoband® (A11158) Images



Western blot analysis of DCHS2 using anti-DCHS2 antibody (A11158). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human U251 whole cell lysates, Lane 2: human U2OS whole cell lysates, Lane 3: human K562 whole cell lysates, Lane 4: rat stomach tissue lysates, Lane 5: rat testis tissue lysates, Lane 6: rat UMR-106 whole cell lysates, Lane 7: mouse testis tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-DCHS2 antigen affinity purified polyclonal antibody (A11158) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for DCHS2 at approximately 90 kDa. The expected band size for DCHS2 is at 370 kDa.

Anti-DCHS2 Antibody

For Research Use Only. Not for use in diagnostic procedures.