

Anti-IDI2 Antibody Picoband® Fluoro647 Conjugated

Catalog Number: A11225-Fluoro647

About IDI2

The protein encoded by this gene catalyzes the conversion of isopentenyl diphosphate to dimethylallyl diphosphate, which is a precursor for the synthesis of cholesterol and other isoprenoids. This gene, which is a product of an ancestral gene duplication event, encodes a protein that may be involved in the aggregation of alpha-synuclein in the cerebral cortex of patients with Lewy body disease. In addition, segmental copy number gains in this locus have been associated with sporadic amyotrophic lateral sclerosis.

Overview

Product Name	Anti-IDI2 Antibody Picoband® Fluoro647 Conjugated
Reactive Species	Human
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, Flow Cytometry, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% Na ₃ N.
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	Q9BXS1

Technical Details

Immunogen	E.coli-derived human IDI2 recombinant protein (Position: M1-V227). Human IDI2 shares 68.2% and 62.6% amino acid (aa) sequence identity with mouse and rat IDI2, respectively.
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	Fluoro647 Excitation Wavelength: 650 nm Emission Wavelength: 665 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-IDI2 Antibody - Fluoro647

For Research Use Only. Not for use in diagnostic procedures.