

## Anti-Centrosomal protein of 83 kDa CEP83 Antibody

Catalog Number: A11554


### Overview

|                      |   |
|----------------------|---|
| Product Name         | Anti-Centrosomal protein of 83 kDa CEP83 Antibody   |
| Reactive Species     | Human, Mouse  |
| Description          | Boster Bio Anti-Centrosomal protein of 83 kDa CEP83 Antibody catalog # A11554. Tested in WB applications. This antibody reacts with Human, Mouse. |
| Application          | WB  |
| Clonality            | Polyclonal  |
| Formulation          | Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2  |
| Storage Instructions | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.        |
| Host                 | Rabbit  |
| Uniprot ID           | Q9Y592  |

### Technical Details

|                     |   |
|---------------------|---|
| Immunogen           | Recombinant fusion protein of human CEP83(NP_057206.2).   |
| Isotype             | IgG   |
| Form                | Liquid  |
| Concentration       | 1 mg/ml   |
| Purification        | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE). |
| Suggested Dilutions | WB: 1:500-1:2000  |

## Anti-Centrosomal protein of 83 kDa CEP83 Antibody (A11554) Images



Western blot analysis of extracts of various cell lines, using CEP83 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Centrosomal protein of 83 kDa CEP83 Antibody

For Research Use Only. Not for use in diagnostic procedures.