

Anti-SAMD8 Antibody Picoband®

Catalog Number: A11828-1

About SAMD8

Predicted to enable ceramide cholinephosphotransferase activity and sphingomyelin synthase activity. Involved in ceramide biosynthetic process and regulation of ceramide biosynthetic process. Located in cytosol and endoplasmic reticulum membrane.


Overview

Product Name	Anti-SAMD8 Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-SAMD8 Antibody Picoband® catalog # A11828-1. Tested in WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q96LT4

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human SAMD8.
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Human

Anti-SAMD8 Antibody Picoband® (A11828-1) Images



Western blot analysis of SAMD8 using anti-SAMD8 antibody (A11828-1). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human U251 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SAMD8 antigen affinity purified polyclonal antibody (A11828-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for SAMD8 at approximately 48 kDa. The expected band size for SAMD8 is at 48 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-SAMD8 Antibody

For Research Use Only. Not for use in diagnostic procedures.