

Anti-PDPR Antibody Picoband®

Catalog Number: A12185

About PDPR

Pyruvate dehydrogenase phosphatase regulatory subunit is a protein that in humans is encoded by the PDPR gene. Pyruvate dehydrogenase complex (PDC) catalyzes the oxidative decarboxylation of pyruvate and links glycolysis to the tricarboxylic acid cycle and fatty acid synthesis. The dephosphorylation and reactivation of PDC is catalyzed by pyruvate dehydrogenase phosphatase (PDP). The dimeric PDP has a catalytic subunit and a regulatory subunit. This gene encodes the FAD-containing regulatory subunit of PDP. The encoded protein acts to decrease the sensitivity of the PDP catalytic subunit to magnesium ions. Alternative splicing results in multiple transcript variants encoding different isoforms.

Overview


Product Name	Anti-PDPR Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-PDPR Antibody Picoband® catalog # A12185. Tested in WB, FCM, ELISA applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q8NCN5

Technical Details


Immunogen	E.coli-derived human PDPR recombinant protein (Position: E155-E655). Human PDPR shares 94.4% amino acid (aa) sequence identity with mouse PDPR.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Mouse, Rat Flow Cytometry (Fixed), 1-3 ug/1x10 ⁶ cells, Human ELISA, 0.1-0.5 ug/ml, -

Anti-PDPR Antibody Picoband® (A12185) Images



Western blot analysis of PDPR using anti-PDPR antibody (A12185). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat heart tissue lysates, Lane 2: mouse heart tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PDPR antigen affinity purified polyclonal antibody (Catalog # A12185) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for PDPR at approximately 99 kDa. The expected band size for PDPR is at 99 kDa.



Flow Cytometry analysis of JK cells using anti-PDPR antibody (A12185). Overlay histogram showing JK cells stained with A12185 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-PDPR Antibody (A12185, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-PDPR Antibody

For Research Use Only. Not for use in diagnostic procedures.