

Anti-SGT2/SGTB Antibody Picoband®

Catalog Number: A13056-1

About SGTB

Predicted to enable molecular adaptor activity. Predicted to be involved in positive regulation of chaperone-mediated protein folding; posttranslational protein targeting to endoplasmic reticulum membrane; and ubiquitin-dependent ERAD pathway. Predicted to be part of TRC complex. Predicted to be active in membrane.

Overview

| | |
|----------------------|--|
| Product Name | Anti-SGT2/SGTB Antibody Picoband® |
| Reactive Species | Human, Rat |
| Description | Boster Bio Anti-SGT2/SGTB Antibody Picoband® catalog # A13056-1. Tested in ELISA, WB applications. This antibody reacts with Human, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance. |
| Application | ELISA, WB |
| Clonality | Polyclonal |
| Formulation | Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ . |
| Storage Instructions | At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing. |
| Host | Rabbit |
| Uniprot ID | Q96EQ0 |


Technical Details

| | |
|-------------------------------|---|
| Immunogen | E.coli-derived human SGT2/SGTB recombinant protein (Position: Q20-A268). |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot. |
| Cross Reactivity | No cross-reactivity with other proteins |
| Isotype | Rabbit IgG |
| Form | Lyophilized |
| Concentration | Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml. |
| Purification | Immunogen affinity purified. |

Suggested Dilutions

Western blot, 0.25-0.5 ug/ml, Human, Rat
ELISA, 0.1-0.5 ug/ml, -

Anti-SGT2/SGTB Antibody Picoband® (A13056-1) Images



Western blot analysis of SGT2/SGTB using anti-SGT2/SGTB antibody (A13056-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates, Lane 2: rat brain tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SGT2/SGTB antigen affinity purified polyclonal antibody (Catalog # A13056-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for SGT2/SGTB at approximately 36 kDa. The expected band size for SGT2/SGTB is at 34 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-SGT2/SGTB Antibody

For Research Use Only. Not for use in diagnostic procedures.