

Anti-SDCCAG10/CWC27 Antibody

Catalog Number: A13346-1

About CWC27

Predicted to enable peptidyl-prolyl cis-trans isomerase activity. Predicted to be involved in protein folding. Located in nucleoplasm. Part of U2-type pre-catalytic spliceosome and catalytic step 2 spliceosome.

Overview

Product Name	Anti-SDCCAG10/CWC27 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-SDCCAG10/CWC27 Antibody catalog # A13346-1. Tested in WB, ICC, IF, IP applications. This antibody reacts with Human, Mouse, Rat.
Application	IP, IF, ICC, WB
Clonality	Polyclonal
Formulation	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg stabilizing protein and 50% glycerol This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	12 months from date of receipt at -20°C as supplied. 6 months at 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q6UX04

Technical Details

Immunogen	E.coli-derived human SDCCAG10/CWC27 recombinant protein (Position: N3-C288).
Form	Liquid
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 1:500-2000 Immunocytochemistry/Immunofluorescence, 1:50-400 ImmunoPrecipitation, 1:250-300

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-SDCCAG10/CWC27 Antibody

For Research Use Only. Not for use in diagnostic procedures.