

Anti-MFSD3 Antibody Picoband™

Catalog Number: A19348-2

About MFSD3

MFSD3 belongs to the major facilitator superfamily. The exact functions of MFSD3 remain unknown.

Overview

Product Name	Anti-MFSD3 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-MFSD3 Antibody Picoband™ catalog # A19348-2. Tested in ELISA, WB, Flow Cytometry applications. This antibody reacts with Human.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal 1B9
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q96ES6

Technical Details

Immunogen	E.coli-derived human MFSD3 recombinant protein (Position: M1-A381). Human MFSD3 shares 79.5% and 78.4% amino acid (aa) sequence identity with mouse and rat MFSD3, respectively.
Predicted Reactive Species	Bovine, Canine, Chicken, Primate, Sheep, Xenopus, Zebrafish
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:

Western blot, 0.25-0.5 µg/ml, Human

Flow Cytometry (Fixed), 1-3 µg/1x10⁶ cells, Human

ELISA, 0.1-0.5 µg/ml, Human

Anti-MFSD3 Antibody Picoband™ (A19348-2) Images




Figure 1. Western blot analysis of MFSD3 using anti-MFSD3 antibody (A19348-2).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human A549 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MFSD3 antigen affinity purified polyclonal antibody (Catalog # A19348-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for MFSD3 at approximately 43 kDa. The expected band size for MFSD3 is at 43 kDa.




Figure 2. Flow Cytometry analysis of 293T cells using anti-MFSD3 antibody (A19348-2).

Overlay histogram showing 293T cells stained with A19348-2 (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-MFSD3 Antibody (A19348-2, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-MFSD3 Antibody