

## Anti-ZCCHC11 TUT4 Antibody

Catalog Number: A32031


### Overview

Product Name	Anti-ZCCHC11 TUT4 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-ZCCHC11 TUT4 Antibody catalog # A32031. Tested in WB,ICC/IF applications. This antibody reacts with Human,Mouse,Rat.
Application	IF, ICC, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q5TAX3


### Technical Details

Immunogen	Recombinant fusion protein of human ZCCHC11(NP_001009881.1).
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).
Suggested Dilutions	WB: 1:500-1:1000 ICC/IF: 1:50-1:200


## Anti-ZCCHC11 TUT4 Antibody (A32031) Images




Western blot analysis of extracts of 293T cells, using ZCCHC11 Antibody at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 20s.



Immunofluorescence analysis of U2OS cells using ZCCHC11 Rabbit pAb at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using ZCCHC11 Rabbit pAb at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of H9C2 cells using ZCCHC11 Rabbit pAb at dilution of 1:100. Blue: DAPI for nuclear staining.

**Submit a product review to Biocompare.com**

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ZCCHC11 TUT4 Antibody

For Research Use Only. Not for use in diagnostic procedures.