

Anti-Zebrafish DTNBA Antibody Picoband®

Catalog Number: AZA0A0R4IAV8

About DTNBA

Predicted to enable zinc ion binding activity. Predicted to be involved in synaptic signaling. Predicted to be located in cytoplasm. Predicted to be active in plasma membrane and synapse. Is expressed in several structures, including brain; hindbrain neural keel; myotome; presumptive diencephalon; and somite border. Orthologous to human DTNB (dystrobrevin beta).


Overview

Product Name	Anti-Zebrafish DTNBA Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish DTNBA Antibody Picoband® catalog # AZA0A0R4IAV8. Tested in WB, IHC applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	A0A0R4IAV8


Technical Details

Immunogen	E.coli-derived Zebrafish DTNBA recombinant protein (Position: A13-Q544).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Zebrafish

Anti-Zebrafish DTNBA Antibody Picoband® (AZA0A0R4IAV8) Images



Western blot analysis of DTNBA using anti-DTNBA antibody (AZA0A0R4IAV8). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish head tissue lysates, Lane 2: whole female zebrafish tissue lysates, Lane 3: whole male zebrafish tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-DTNBA antigen affinity purified polyclonal antibody (AZA0A0R4IAV8) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for DTNBA at approximately 66 kDa. The expected band size for DTNBA is at 66 kDa.



IHC analysis of DTNBA using anti-DTNBA antibody (AZA0A0R4IAV8). DTNBA was detected in a paraffin-embedded section of zebrafish spinal cord tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-DTNBA Antibody (AZA0A0R4IAV8) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Zebrafish DTNBA Antibody

For Research Use Only. Not for use in diagnostic procedures.