

Anti-Zebrafish DRD4A Antibody Picoband®

Catalog Number: AZA0A0R4IR92

About DRD4A

Predicted to enable G protein-coupled receptor activity and dopamine neurotransmitter receptor activity, coupled via Gi/Go. Acts upstream of or within spinal cord motor neuron cell fate specification and ventral spinal cord interneuron differentiation. Predicted to be located in membrane. Predicted to be active in plasma membrane. Is expressed in nervous system. Human ortholog(s) of this gene implicated in several diseases, including Gilles de la Tourette syndrome; antisocial personality disorder; attention deficit hyperactivity disorder; conduct disorder; and substance abuse (multiple). Orthologous to human DRD4 (dopamine receptor D4).


Overview

Product Name	Anti-Zebrafish DRD4A Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish DRD4A Antibody Picoband® catalog # AZA0A0R4IR92. Tested in WB applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	A0A0R4IR92

Technical Details

Immunogen	E.coli-derived Zebrafish DRD4A recombinant protein (Position: M1-R300).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish

Anti-Zebrafish DRD4A Antibody Picoband® (AZA0A0R4IR92) Images



Western blot analysis of DRD4A using anti-DRD4A antibody (AZA0A0R4IR92). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish head tissue lysates, Lane 2: whole female zebrafish tissue lysates, Lane 3: whole male zebrafish tissue lysates, Lane 4: zebrafish embryo tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-DRD4A antigen affinity purified polyclonal antibody (AZA0A0R4IR92) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for DRD4A at approximately 43 kDa. The expected band size for DRD4A is at 43 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Zebrafish DRD4A Antibody

For Research Use Only. Not for use in diagnostic procedures.