

Anti-Zebrafish FDFT1 Antibody Picoband®

Catalog Number: AZA0A0R4IVV5

About FDFT1

Predicted to enable squalene synthase [NAD(P)H] activity. Predicted to be involved in cholesterol biosynthetic process and farnesyl diphosphate metabolic process. Predicted to act upstream of or within lipid biosynthetic process. Predicted to be located in membrane. Predicted to be active in endoplasmic reticulum membrane. Is expressed in brain. Orthologous to human FDFT1 (farnesyl-diphosphate farnesyltransferase 1).

Submit a product review to Biocompare.com

[Submit a review of this product](#) to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Overview


Product Name	Anti-Zebrafish FDFT1 Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish FDFT1 Antibody Picoband® catalog # AZA0A0R4IVV5. Tested in WB applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	A0A0R4IVV5

Technical Details

Immunogen	E.coli-derived Zebrafish FDFT1 recombinant protein (Position: R25-E373).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish

Anti-Zebrafish FDFT1 Antibody Picoband® (AZA0A0R4IVV5) Images



Western blot analysis of FDFT1 using anti-FDFT1 antibody (AZA0A0R4IVV5). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish head tissue lysates, Lane 2: whole female zebrafish tissue lysates, Lane 3: whole male zebrafish tissue lysates, Lane 4: zebrafish embryo tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-FDFT1 antigen affinity purified polyclonal antibody (AZA0A0R4IVV5) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for FDFT1 at approximately 50 kDa. The expected band size for FDFT1 is at 47 kDa.

[Anti-Zebrafish FDFT1 Antibody](#)

For Research Use Only. Not for use in diagnostic procedures.