

Anti-Zebrafish HEG1 Antibody Picoband®

Catalog Number: AZA9C3V5

About HEG1

Predicted to enable calcium ion binding activity. Acts upstream of or within cell-cell junction assembly and heart development. Predicted to be located in anchoring junction; extracellular region; and plasma membrane. Is expressed in cardiovascular system; central nervous system; somite; and ventral mesoderm. Orthologous to human HEG1 (heart development protein with EGF like domains 1).

Submit a product review to Biocompare.com

[Submit a review of this product](#) to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Overview


Product Name	Anti-Zebrafish HEG1 Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish HEG1 Antibody Picoband® catalog # AZA9C3V5. Tested in WB applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	A9C3V5

Technical Details

Immunogen	E.coli-derived Zebrafish HEG1 recombinant protein (Position: E236-L492).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish

Anti-Zebrafish HEG1 Antibody Picoband® (AZA9C3V5) Images



Western blot analysis of HEG1 using anti-HEG1 antibody (AZA9C3V5). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: whole female zebrafish tissue lysates, Lane 2: whole male zebrafish tissue lysates, Lane 3: zebrafish embryo tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-HEG1 antigen affinity purified polyclonal antibody (AZA9C3V5) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for HEG1 at approximately 105 kDa. The expected band size for HEG1 is at 105 kDa.

Anti-Zebrafish HEG1 Antibody

For Research Use Only. Not for use in diagnostic procedures.