BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Zebrafish RECQL4 Antibody Picoband®

Catalog Number: AZE7F2M7

About RECQL4

Predicted to enable 3'-5' DNA helicase activity and four-way junction helicase activity. Predicted to be involved in DNA replication; double-strand break repair via homologous recombination; and telomere maintenance. Predicted to be active in chromosome; cytoplasm; and nucleus. Human ortholog(s) of this gene implicated in Baller-Gerold syndrome; Rothmund-Thomson syndrome; and rapadilino syndrome. Orthologous to human RECQL4 (RecQ like helicase 4).

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-Zebrafish RECQL4 Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish RECQL4 Antibody Picoband® catalog # AZE7F2M7. Tested in WB applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	E7F2M7

Technical Details


Immunogen	E.coli-derived Zebrafish RECQL4 recombinant protein (Position: H81-R1459).
Form	Lyophilized

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Zebrafish RECQL4 Antibody Picoband® (AZE7F2M7) Images



Western blot analysis of RECQL4 using anti-RECQL4 antibody (AZE7F2M7). Electrophoresis was performed on a 8% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: whole female zebrafish lysates, Lane 2: whole male zebrafish lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-RECQL4 antigen affinity purified polyclonal antibody (AZE7F2M7) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for RECQL4 at approximately 163 kDa. The expected band size for RECQL4 is at 163 kDa.

Anti-Zebrafish RECQL4 Antibody

For Research Use Only. Not for use in diagnostic procedures.