

Anti-Zebrafish HSD3B1 Antibody Picoband®

Catalog Number: AZF1QSA2

About HSD3B1

Enables 3-beta-hydroxy-Delta5-steroid dehydrogenase (NAD+) activity. Acts upstream of or within pigmentation. Is expressed in brain; endocrine system; and gonad primordium. Human ortholog(s) of this gene implicated in hypertension. Orthologous to human HSD3B1 (hydroxy-delta-5-steroid dehydrogenase, 3 beta- and steroid delta-isomerase 1).

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.




Overview

Product Name	Anti-Zebrafish HSD3B1 Antibody Picoband®
Reactive Species	Zebrafish
Description	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. Immunogen affinity purified. Enables 3-beta-hydroxy-Delta5-steroid dehydrogenase (NAD+) activity. Acts upstream of or within pigmentation. Is expressed in brain; endocrine system; and gonad primordium. Human ortholog(s) of this gene implicated in hypertension. Orthologous to human HSD3B1 (hydroxy-delta-5-steroid dehydrogenase, 3 beta- and steroid delta-isomerase 1). E.coli-derived Zebrafish HSD3B1 recombinant protein (Position: F16-K374) anti-zebrafish-hsd3b1-antibody-azf1qsa2-boster Anti-Zebrafish HSD3B1 Antibody Picoband® Boster Bio Anti-Zebrafish HSD3B1 Antibody Picoband® catalog # AZF1QSA2. Tested in WB, IHC applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	F1QSA2


Technical Details

Immunogen	E.coli-derived Zebrafish HSD3B1 recombinant protein (Position: F16-K374)
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Zebrafish


Anti-Zebrafish HSD3B1 Antibody Picoband® (AZF1QSA2) Images



Western blot analysis of HSD3B1 using anti-HSD3B1 antibody (AZF1QSA2). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish head tissue lysates, Lane 2: whole female zebrafish tissue lysates, Lane 3: whole male zebrafish tissue lysates, Lane 4: zebrafish embryo tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-HSD3B1 antigen affinity purified polyclonal antibody (AZF1QSA2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for HSD3B1 at approximately 41 kDa. The expected band size for HSD3B1 is at 41 kDa.



IHC analysis of HSD3B1 using anti-HSD3B1 antibody (AZF1QSA2). HSD3B1 was detected in a paraffin-embedded section of zebrafish head kidney tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-HSD3B1 Antibody (AZF1QSA2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of HSD3B1 using anti-HSD3B1 antibody (AZF1QSA2). HSD3B1 was detected in a paraffin-embedded section of zebrafish brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-HSD3B1 Antibody (AZF1QSA2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

For Research Use Only. Not for use in diagnostic procedures.