BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Zebrafish NPPA Antibody Picoband®

Catalog Number: AZF1QVW0

About NPPA

Predicted to enable hormone activity and hormone receptor binding activity. Acts upstream of or within atrioventricular canal development; cardiac jelly development; and cardiac muscle cell proliferation. Predicted to be located in extracellular region. Predicted to be active in cytoplasm and extracellular space. Is expressed in heart rudiment; pericardial region; and skeletal muscle cell. Human ortholog(s) of this gene implicated in artery disease (multiple); heart conduction disease (multiple); and peritonitis. Orthologous to human NPPA (natriuretic peptide A).

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-Zebrafish NPPA Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish NPPA Antibody Picoband® catalog # AZF1QVW0. Tested in WB applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	F1QVW0

Technical Details

Immunogen	E.coli-derived Zebrafish NPPA recombinant protein (Position: H23-G139).
Form	Lyophilized

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Zebrafish NPPA Antibody Picoband® (AZF1QVW0) Images



Western blot analysis of NPPA using anti-NPPA antibody (AZF1QVW0). Electrophoresis was performed on a 12% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish head tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-NPPA antigen affinity purified polyclonal antibody (AZF1QVW0) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for NPPA at approximately 15 kDa. The expected band size for NPPA is at 15 kDa.

Anti-Zebrafish NPPA Antibody

For Research Use Only. Not for use in diagnostic procedures.