

Anti-Zebrafish IGF2BP1 Antibody Picoband®

Catalog Number: AZQ08CK7

About IGF2BP1

Predicted to enable N6-methyladenosine-containing RNA reader activity and mRNA 3'-UTR binding activity. Acts upstream of or within neural retina development and regulation of hepatocyte proliferation. Predicted to be located in P-body and cytoplasmic stress granule. Predicted to be part of CRD-mediated mRNA stability complex. Predicted to be active in cytosol and nucleus. Is expressed in brain; digestive system; and head. Orthologous to human IGF2BP1 (insulin like growth factor 2 mRNA binding protein 1).

Submit a product review to Biocompare.com

[Submit a review of this product](#) to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Overview


Product Name	Anti-Zebrafish IGF2BP1 Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish IGF2BP1 Antibody Picoband® catalog # AZQ08CK7. Tested in WB applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q08CK7

Technical Details

Immunogen	E.coli-derived Zebrafish IGF2BP1 recombinant protein (Position: N8-K598).
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish

Anti-Zebrafish IGF2BP1 Antibody Picoband® (AZQ08CK7) Images



Western blot analysis of IGF2BP1 using anti-IGF2BP1 antibody (AZQ08CK7). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish embryo tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-IGF2BP1 antigen affinity purified polyclonal antibody (AZQ08CK7) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for IGF2BP1 at approximately 70 kDa. The expected band size for IGF2BP1 is at 65 kDa.

Anti-Zebrafish IGF2BP1 Antibody

For Research Use Only. Not for use in diagnostic procedures.