

Anti-Zebrafish MYPT1/PPP1R12A Antibody Picoband®

Catalog Number: AZQ6DRG7

About PPP1R12A

PPP1R12A (Protein phosphatase 1 regulatory subunit 12A), also called MYPT1 (Myosin phosphatase target subunit 1), is an enzyme that in humans is encoded by the PPP1R12A gene. PPP1R12A is one of the subunits of myosin phosphatase. Sequencing analysis showed that human PPP1R12A contains 1,030 amino acids with a calculated molecular mass of approximately 115 kD. The PPP1R12A gene is mapped on 12q21.2-q21.3. PPP1R12A is the protein that regulates PP1 function in smooth muscle relaxation. The cellular MYPT1-PP1-delta -specific inhibitor CPI17 caused a loss of merlin function characterized by merlin phosphorylation, Ras activation, and transformation. Jin et al. concluded that PPP1R12A and its substrate merlin are part of a previously undescribed tumor suppressor cascade that can be hindered in two ways, by mutation of the NF2 gene and by upregulation of the oncoprotein CPI17.

Overview


Product Name	Anti-Zebrafish MYPT1/PPP1R12A Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish MYPT1/PPP1R12A Antibody Picoband® catalog # AZQ6DRG7. Tested in WB, IHC applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q6DRG7

Technical Details


Immunogen	E.coli-derived zebrafish MYPT1/PPP1R12A recombinant protein (Position: M1-D40).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish

Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Zebrafish


Anti-Zebrafish MYPT1/PPP1R12A Antibody Picoband® (AZQ6DRG7) Images



Western blot analysis of PPP1R12A using anti-PPP1R12A antibody (AZQ6DRG7). Electrophoresis was performed on a 8% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: whole female zebrafish tissue lysates, Lane 2: whole male zebrafish tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PPP1R12A antigen affinity purified polyclonal antibody (AZQ6DRG7) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for PPP1R12A at approximately 140 kDa. The expected band size for PPP1R12A is at 115 kDa.




IHC analysis of PPP1R12A using anti-PPP1R12A antibody (AZQ6DRG7). PPP1R12A was detected in a paraffin-embedded section of zebrafish brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-PPP1R12A Antibody (AZQ6DRG7) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of PPP1R12A using anti-PPP1R12A antibody (AZQ6DRG7). PPP1R12A was detected in a paraffin-embedded section of zebrafish liver tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-PPP1R12A Antibody (AZQ6DRG7) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

IHC analysis of PPP1R12A using anti-PPP1R12A antibody (AZQ6DRG7). PPP1R12A was detected in a paraffin-embedded section of zebrafish muscle tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked



with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-PPP1R12A Antibody (AZQ6DRG7) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Zebrafish MYPT1/PPP1R12A Antibody

For Research Use Only. Not for use in diagnostic procedures.