

Anti-Zebrafish DPYD Antibody Picoband®

Catalog Number: AZQ6NYG8

About DPYD

Predicted to enable NADP binding activity; dihydropyrimidine dehydrogenase (NADP+) activity; and uracil binding activity. Predicted to be involved in thymidine catabolic process; thymine catabolic process; and uracil catabolic process. Predicted to be located in cytoplasm. Predicted to be active in cytosol. Is expressed in intestinal bulb; liver; pronephric duct; and yolk syncytial layer. Human ortholog(s) of this gene implicated in acute lymphoblastic leukemia; dihydropyrimidine dehydrogenase deficiency; and pancreatic cancer. Orthologous to human DPYD (dihydropyrimidine dehydrogenase).


Overview

Product Name	Anti-Zebrafish DPYD Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish DPYD Antibody Picoband® catalog # AZQ6NYG8. Tested in WB, IHC applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q6NYG8

Technical Details

Immunogen	E.coli-derived Zebrafish DPYD recombinant protein (Position: K35-N954).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Zebrafish

Anti-Zebrafish DPYD Antibody Picoband® (AZQ6NYG8) Images



Western blot analysis of DPYD using anti-DPYD antibody (AZQ6NYG8). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 4: zebrafish embryo tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-DPYD antigen affinity purified polyclonal antibody (AZQ6NYG8) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for DPYD at approximately 111 kDa. The expected band size for DPYD is at 111 kDa.



IHC analysis of DPYD using anti-DPYD antibody (AZQ6NYG8). DPYD was detected in a paraffin-embedded section of zebrafish eye tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-DPYD Antibody (AZQ6NYG8) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Zebrafish DPYD Antibody

For Research Use Only. Not for use in diagnostic procedures.