

Anti-Zebrafish DGAT1A Antibody Picoband®

Catalog Number: AZQ6P3J0

About DGAT1A

Predicted to enable diacylglycerol O-acyltransferase activity. Acts upstream of or within regulation of response to tumor cell. Predicted to be located in endoplasmic reticulum and membrane. Predicted to be active in endoplasmic reticulum membrane. Is expressed in gut; liver; and muscle. Human ortholog(s) of this gene implicated in cardiomyopathy and congenital diarrhea 7 with exudative enteropathy. Orthologous to human DGAT1 (diacylglycerol O-acyltransferase 1).


Overview

Product Name	Anti-Zebrafish DGAT1A Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish DGAT1A Antibody Picoband® catalog # AZQ6P3J0. Tested in WB applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q6P3J0

Technical Details

Immunogen	E.coli-derived Zebrafish DGAT1A recombinant protein (Position: E21-D372).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish

Anti-Zebrafish DGAT1A Antibody Picoband® (AZQ6P3J0) Images



Western blot analysis of DGAT1A using anti-DGAT1A antibody (AZQ6P3J0). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish head tissue lysates, Lane 2: whole female zebrafish tissue lysates, Lane 3: whole male zebrafish tissue lysates, Lane 4: zebrafish embryo tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-DGAT1A antigen affinity purified polyclonal antibody (AZQ6P3J0) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for DGAT1A at approximately 50 kDa. The expected band size for DGAT1A is at 57 kDa.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Zebrafish DGAT1A Antibody

For Research Use Only. Not for use in diagnostic procedures.