

## Anti-Zebrafish FSHB Antibody Picoband®

Catalog Number: AZQ6TCF7

### About FSHB

Enables hormone activity. Acts upstream of or within ovarian follicle development; sex determination; and spermatogenesis. Predicted to be located in extracellular region. Predicted to be active in cytoplasm and extracellular space. Is expressed in several structures, including brain; endocrine system; female organism; kidney; and male organism. Human ortholog(s) of this gene implicated in amenorrhea; hypogonadotropic hypogonadism 23 with or without anosmia; and hypogonadotropic hypogonadism 24 without anosmia. Orthologous to several human genes including CGB1 (chorionic gonadotropin subunit beta 1); CGB2 (chorionic gonadotropin subunit beta 2); and CGB3 (chorionic gonadotropin subunit beta 3).


### Overview

Product Name	Anti-Zebrafish FSHB Antibody Picoband®
Reactive Species	Zebrafish
Description	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. Immunogen affinity purified. Enables hormone activity. Acts upstream of or within ovarian follicle development; sex determination; and spermatogenesis. Predicted to be located in extracellular region. Predicted to be active in cytoplasm and extracellular space. Is expressed in several structures, including brain; endocrine system; female organism; kidney; and male organism. Human ortholog(s) of this gene implicated in amenorrhea; hypogonadotropic hypogonadism 23 with or without anosmia; and hypogonadotropic hypogonadism 24 without anosmia. Orthologous to several human genes including CGB1 (chorionic gonadotropin subunit beta 1); CGB2 (chorionic gonadotropin subunit beta 2); and CGB3 (chorionic gonadotropin subunit beta 3). E.coli-derived Zebrafish FSHB recombinant protein (Position: E19-H130) anti-zebrafish-fshb-antibody-azq6tcf7-boster. Anti-Zebrafish FSHB Antibody Picoband® Boster Bio Anti-Zebrafish FSHB Antibody Picoband® catalog # AZQ6TCF7. Tested in WB, IHC applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q6TCF7


### Technical Details

Immunogen	E.coli-derived Zebrafish FSHB recombinant protein (Position: E19-H130)
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Zebrafish


## Anti-Zebrafish FSHB Antibody Picoband® (AZQ6TCF7) Images



Western blot analysis of FSHB using anti-FSHB antibody (AZQ6TCF7). Electrophoresis was performed on a 12% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish head tissue lysates, Lane 2: zebrafish embryo tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-FSHB antigen affinity purified polyclonal antibody (AZQ6TCF7) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for FSHB at approximately 20 kDa. The expected band size for FSHB is at 14 kDa.




IHC analysis of FSHB using anti-FSHB antibody (AZQ6TCF7). FSHB was detected in a paraffin-embedded section of zebrafish ovary tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-FSHB Antibody (AZQ6TCF7) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




IHC analysis of FSHB using anti-FSHB antibody (AZQ6TCF7). FSHB was detected in a paraffin-embedded section of zebrafish testis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-FSHB Antibody (AZQ6TCF7) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

IHC analysis of FSHB using anti-FSHB antibody (AZQ6TCF7). FSHB was detected in a paraffin-embedded section of zebrafish testis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-FSHB Antibody (AZQ6TCF7) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



rabbit anti-FSHB Antibody (AZQ6TCF7) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of FSHB using anti-FSHB antibody (AZQ6TCF7). FSHB was detected in a paraffin-embedded section of zebrafish eye tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-FSHB Antibody (AZQ6TCF7) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-Zebrafish FSHB Antibody

For Research Use Only. Not for use in diagnostic procedures.