

Anti-Zebrafish DICER1 Antibody Picoband®

Catalog Number: AZQ6TV19

About DICER1

Dicer(DICER1), also known as endoribonuclease Dicer or helicase with RNase motif, is an enzyme that in humans is encoded by the DICER1 gene. It is mapped to 14q32.13. The DICER1 gene, a member of the ribonuclease III (RNaseIII) family, is involved in the generation of microRNAs (miRNAs), which modulate gene expression at the posttranscriptional level. DICER1 possesses an RNA helicase motif containing a DEXH box in its amino terminus and an RNA motif in the carboxy terminus. DICER, also known as helicase-MOI, is required by the RNA interference and small temporal RNA (stRNA) pathways to produce the active small RNA component that represses gene expression. In addition, DICER1 is required for formation of the RNA induced silencing complex (RISC). It also cleaves double-stranded RNA to produce short interfering RNAs (siRNAs) which target the selective destruction of complementary RNAs.

Overview


Product Name	Anti-Zebrafish DICER1 Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish-DICER1-Antibody Picoband® catalog # AZQ6TV19. Tested in WB applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q6TV19

Technical Details

Immunogen	E.coli-derived zebrafish DICER1 recombinant protein (Position: Y1522-N1865).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish



Anti-Zebrafish DICER1 Antibody Picoband® (AZQ6TV19) Images



Western blot analysis of DICER1 using anti-DICER1 antibody (AZQ6TV19). Electrophoresis was performed on a 8% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish embryo tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-DICER1 antigen affinity purified polyclonal antibody (AZQ6TV19) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for DICER1 at approximately 250 kDa. The expected band size for DICER1 is at 219 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Zebrafish DICER1 Antibody

For Research Use Only. Not for use in diagnostic procedures.