

Anti-Zebrafish DDIT4 Antibody Picoband®

Catalog Number: AZQ7T346

About DDIT4

Predicted to enable 14-3-3 protein binding activity. Acts upstream of or within dorsal/ventral pattern formation. Predicted to be located in cytoplasm. Is expressed in several structures, including central nervous system; digestive system; mesoderm; muscle; and pleuroperitoneal region. Orthologous to human DDIT4 (DNA damage inducible transcript 4).


Overview

Product Name	Anti-Zebrafish DDIT4 Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish DDIT4 Antibody Picoband® catalog # AZQ7T346. Tested in WB applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q7T346

Technical Details

Immunogen	E.coli-derived Zebrafish DDIT4 recombinant protein (Position: M1-C220).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish

Anti-Zebrafish DDIT4 Antibody Picoband® (AZQ7T346) Images



Western blot analysis of DDIT4 using anti-DDIT4 antibody (AZQ7T346). Electrophoresis was performed on a 12% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish embryo tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-DDIT4 antigen affinity purified polyclonal antibody (AZQ7T346) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for DDIT4 at approximately 24 kDa. The expected band size for DDIT4 is at 24 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Zebrafish DDIT4 Antibody

For Research Use Only. Not for use in diagnostic procedures.