

## Anti-Zebrafish VANGL2 Antibody Picoband®

Catalog Number: AZQ8UVJ6

### About VANGL2

Involved in several processes, including Wnt signaling pathway, planar cell polarity pathway; nervous system development; and regulation of Wnt signaling pathway. Acts upstream of or within several processes, including cytoskeleton organization; embryonic morphogenesis; and generation of neurons. Located in plasma membrane. Is expressed in central nervous system; liver; male organism; pharyngeal arch; and sensory system. Used to study ciliopathy. Human ortholog(s) of this gene implicated in neural tube defect. Orthologous to human VANGL2 (VANGL planar cell polarity protein 2).

### Overview


|                      |   |
|----------------------|---|
| Product Name         | Anti-Zebrafish VANGL2 Antibody Picoband®  |
| Reactive Species     | Zebrafish   |
| Description          | Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. Immunogen affinity purified. Involved in several processes, including Wnt signaling pathway, planar cell polarity pathway; nervous system development; and regulation of Wnt signaling pathway. Acts upstream of or within several processes, including cytoskeleton organization; embryonic morphogenesis; and generation of neurons. Located in plasma membrane. Is expressed in central nervous system; liver; male organism; pharyngeal arch; and sensory system. Used to study ciliopathy. Human ortholog(s) of this gene implicated in neural tube defect. Orthologous to human VANGL2 (VANGL planar cell polarity protein 2). E.coli-derived Zebrafish VANGL2 recombinant protein (Position: M1-D489) anti-zebrafish-vangl2-antibody-azq8uvj6-boster Anti-Zebrafish VANGL2 Antibody Picoband® Boster Bio Anti-Zebrafish VANGL2 Antibody Picoband® catalog # AZQ8UVJ6. Tested in WB, IHC applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance. |
| Application          | IHC, WB   |
| Clonality            | Polyclonal  |
| Formulation          | Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .   |
| Storage Instructions | At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.   |
| Host                 | Rabbit  |
| Uniprot ID           | Q8UVJ6  |

### Technical Details

|           |   |
|-----------|---|
| Immunogen | E.coli-derived Zebrafish VANGL2 recombinant protein (Position: M1-D489) |
|-----------|---|

|                     |  |
|---------------------|--|
| Form                | Lyophilized  |
| Concentration       | Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.  |
| Purification        | Immunogen affinity purified.   |
| Suggested Dilutions | Western blot, 0.25-0.5 ug/ml, Zebrafish<br>Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Zebrafish |


## Anti-Zebrafish VANGL2 Antibody Picoband® (AZQ8UVJ6) Images



Western blot analysis of VANGL2 using anti-VANGL2 antibody (AZQ8UVJ6). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish head tissue lysates, Lane 2: whole female zebrafish tissue lysates, Lane 3: whole male zebrafish tissue lysates, Lane 4: zebrafish embryo tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-VANGL2 antigen affinity purified polyclonal antibody (AZQ8UVJ6) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for VANGL2 at approximately 60 kDa. The expected band size for VANGL2 is at 60 kDa.



IHC analysis of VANGL2 using anti-VANGL2 antibody (AZQ8UVJ6). VANGL2 was detected in a paraffin-embedded section of zebrafish skin tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-VANGL2 Antibody (AZQ8UVJ6) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of VANGL2 using anti-VANGL2 antibody (AZQ8UVJ6). VANGL2 was detected in a paraffin-embedded section of zebrafish colon tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-VANGL2 Antibody (AZQ8UVJ6) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your



reviews help your fellow scientists make the right decisions. Thank you for your contribution.

**Anti-Zebrafish VANGL2 Antibody**

**For Research Use Only. Not for use in diagnostic procedures.**