

## Anti-Zebrafish GNAT1 Antibody Picoband®

Catalog Number: AZQ90WX6

### About GNAT1

Predicted to enable G protein-coupled receptor binding activity; G-protein beta/gamma-subunit complex binding activity; and GTPase activity. Predicted to be involved in adenylate cyclase-modulating G protein-coupled receptor signaling pathway; detection of light stimulus involved in visual perception; and phototransduction, visible light. Predicted to act upstream of or within G protein-coupled receptor signaling pathway and visual perception. Predicted to be located in membrane. Predicted to be part of heterotrimeric G-protein complex. Predicted to be active in cytoplasm. Is expressed in head; pineal complex; and visual system. Human ortholog(s) of this gene implicated in congenital stationary night blindness 1G; congenital stationary night blindness autosomal dominant 3; and night blindness. Orthologous to human GNAT1 (G protein subunit alpha transducin 1).

### Overview


Product Name	Anti-Zebrafish GNAT1 Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish GNAT1 Antibody Picoband® catalog # AZQ90WX6. Tested in WB, IHC applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q90WX6

### Technical Details


Immunogen	E.coli-derived Zebrafish GNAT1 recombinant protein (Position: M1-E314).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish

Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Zebrafish


## Anti-Zebrafish GNAT1 Antibody Picoband® (AZQ90WX6) Images



Western blot analysis of GNAT1 using anti-GNAT1 antibody (AZQ90WX6). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish head tissue lysates, Lane 2: whole female zebrafish tissue lysates, Lane 3: whole male zebrafish tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-GNAT1 antigen affinity purified polyclonal antibody (AZQ90WX6) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for GNAT1 at approximately 39 kDa. The expected band size for GNAT1 is at 39 kDa.



IHC analysis of GNAT1 using anti-GNAT1 antibody (AZQ90WX6). GNAT1 was detected in a paraffin-embedded section of zebrafish eye tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-GNAT1 Antibody (AZQ90WX6) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of GNAT1 using anti-GNAT1 antibody (AZQ90WX6). GNAT1 was detected in a paraffin-embedded section of zebrafish brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-GNAT1 Antibody (AZQ90WX6) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Zebrafish GNAT1 Antibody

For Research Use Only. Not for use in diagnostic procedures.